

# Care Sheet for the Jackson's Chameleon *Trioceros jacksonii xantholophus*

By Petr Necas & Bill Strand



Legend	Sub-legend	Description
Taxonomy	Taxon	<i>Trioceros jacksonii xantholophus</i>
	Common Names	Yellow-Crested Jackson's Chameleon (English) Three-Horned Chameleon (English) Kiumbu (Meru language) Murtyu (Kikuyu language)
	Original name	<i>Chamaeleo jacksonii xantholophus</i>
	Author	Eason, Ferguson & Hebrard, 1988
	Original description	Copeia, 1988: 586; fig. 1
	Natural Location	3 km south of Meru, Meru district, Kenya
	Typus	NMK L/1796
	Taxonomy	The largest subspecies of the Jackson's Three-Horned Chameleon, <i>Trioceros jacksonii</i> Member of the genus <i>Trioceros</i>
Life Space	Range	Kenya (Eastern slopes of Mt. Kenya, Nyambene Range), Established in Hawaii, USA; Introduced in continental USA (Florida, California)
	Altitude	600 – 1800m a.s.l., typically 1500-1700m
	Macro-habitat	Montane forest, farmland
	Micro-habitat	Forest edge, shrubs, living fences, agricultural plants (Coffee, Tea, Miraa/Khat, Macadamia...)
	Perching Height	0–12m above ground
	Daily Activity	Whole day in the bush or on its edge, morning and late afternoon 1-2 hour basking on a sun exposed branch, sleeping in the middle of bushes, in case of rain hiding in the middle of bushes
	IUCN Status	Least Concern due to widespread natural occurrence and tolerance of habitat modification
	Conservation	Meru forest reserve, Nyambene forest reserve, Maua Forest reserve
	CITES	CITES Ap. II

Legend	Sub-legend	Description
Climate	Climate Type	Equatorial tropical montane climate
	Dry seasons	December to March and June to September
	Rainy seasons	March to May and October to December
	Temperature	Day: Up to 90°F (32°C) with 72-77°F (22-25°C) in the shade Night: 50 - 60°F (10-16°C)
	Humidity	Up to 100 % at night all year long, below 50% (to below 30% in dry season) at daytime
Life Cycle	Parity	Viviparous – live-bearing
	Gestation period	6-7 months
	Clutch size	7-51
	Size at birth	Approx. 2in
	Maturity reached	at 9-12 months
	Mating period	All year around with peak of the births in rainy seasons
	Longevity	2-3 years in the wild, in the captivity 5-6, up to 9 years
Morphology	General	A typical chameleon of middle size, equipped with all typical chameleon features like zygodactylous feet, prehensile tail, independently moving eyes in lid turrets, long prehensile tail, skin capable of color change, The head with typical rough crests consisting of heavily enlarged warty or pointed scales. Possessing a temporal gland used for prey luring and anti-predatory mechanisms.
	Size	Males usually 10-12” up to 16” (25-30cm, up to 42cm), females up to 13” ( 32cm)
	Sexual dimorphism	3 cranial true horns in males (one rostral, 2 preorbital), females hornless often with tiny conical scale on the analogous place of the base of male horns, rarely one short true rostral horn Males, when excited, lime yellow with dark blue casque and light blue tail, females typically uniformly green or with diverse patterns of brown, maroon, gray to black Males have a swollen tail base

Legend	Sub-legend	Description
Health issues	Internal parasites	Roundworms Tapeworms Flukes Treatment: Consult veterinarian
	RI (Respiratory infection)	Symptoms: heavy breathing, visible ribs, gaping, sitting with head up, bubbles in throat Cause: arise often in captivity as a result of too high humidity at daytime combined with high temperatures Treatment: Antibiotics to be prescribed by a veterinarian
	TGI (Temporal Gland infection)	Symptoms: Swollen temporal gland area Cause: arise rarely in captivity as a result of too high humidity at daytime combined with high temperatures Treatment: Antibiotics to be prescribed by a veterinarian
	Mouthrot (Stomatitis; Gingivitis ulcerosa)	Symptoms: white cheese-like deposits along the jaws, swollen jaws Cause: arise often in captivity as a result of injuries of jaws and mucous in combination with husbandry issues Treatment: Antibiotics to be prescribed by a veterinarian
	MBD (Metabolic Bone Disease)	Symptoms: casque and head deformities, rubber jaw, broken bones of extremities, fractures of ribs Cause: A captive condition resulting from an imbalance of vitamin D3 supplementation, lack of Calcium + magnesium in food and/or insufficient UVB exposure Treatment: proper diet and UVB exposure. In heavy cases - veterinarian
	Obesity	Symptoms: Heavy body, inactivity, swollen cheeks and casques, puffy extremities Behavior: Picky eating, slower moving, puffy belly resting on or overflowing branch Cause: Overfeeding Treatment: reduce diet
	Edema (Oedema)	Symptoms: Swelling or puffiness especially around the neck region Cause: imbalance in vitamins Treatment: reduce supplementation
	Eye infections	Symptoms: Swollen turrets or their parts, closed eyes, eye opening blocked by transparent or milky pus Cause: Under/overdose of vitamins or physical trauma Treatment: Antibiotics to be prescribed by a veterinarian



Legend	Sub-legend	Description
<b>Caging</b>	Caging type	<p>Individual</p> <p>Cohabiting of adults not recommended</p> <p>Cohabiting of young juveniles up to 2 months of age possible in densely planted cages with close observation</p> <p>If kept individually, visual contact of at least 3m from each other simulates a natural interaction</p>
	Cage conditions	<p>Day Temperatures: 68-72°F (20-23°C) with basking spot up to 83°F/ 28°C,</p> <p>Night Temperatures: under 50-59° (15°C) and lower in acclimated animals</p> <p>Humidity levels: nighttime up to 100% towards morning, daytime under 50%</p> <p>UVI: 5-7 at basking spot</p>
	Cage size	24 x 24 x 48in or greater for both sexes
	Cage type	Full screen cage or Glass terrariums with ventilation
	Cage interior	<p>Dense foliage from live plants with extensive network of thin natural branches</p> <p>Freely exposed horizontal branch for basking in safe distance (head and body length from the heat source)</p> <p>A thick diagonal rough barked branch in the lower part of cage for birthing</p>
	Lighting	<p>Light bulb white light = 12 hours per day</p> <p>Heat bulb white light (not red) = according to surrounding temperatures</p> <p>Linear UVB bulb = 12 hours per day</p> <p>Nighttime: No heat/light source - including blue and red bulbs!</p>
	Water management	<p>Fog (ultra-sonic humidifier) at night (from 1AM till dawn)</p> <p>Morning Mister: Short misting session (2 minutes) 30 minutes before lights go ON;</p> <p>Purpose is to create a layer of dew on leaves for chameleon to find when it wakes up.</p> <p>Afternoon Rain Shower: During rainy season simulate rain shower by switching off heat lamps for 30 minutes and then run misters a couple minutes; Do not bring heat lamp back on for 30 minutes after shower is over.</p> <p>Evening Misting: Wait until all lights are off and chameleon has settled in. Run mister for two minutes to raise humidity.</p> <p>Dripper: best in the morning hours</p> <p>Use cool or ambient temperature water. Do NOT heat or warm water.</p> <p>Do NOT “bathe” or “soak” your chameleon.</p> <p>Provide dense plant cover so chameleon can choose to get in or out of misting action.</p>

Legend	Sub-legend	Description
Feeding in captivity	Food	<p>General rule: as variable as possible</p> <p>Overfeeding risk: Usually not an issue, but possible. Consult “Obesity” in health section for symptoms.</p> <p>Food items size: preferable smaller size under 1/2 inch</p> <p>To feed: flies, crickets, roaches, superworms, hornworms, fly larvae, wax worms and wax moths, silkworms and silk moths</p> <p>Food to consider: wild bees, small snails</p>
	Supplements	<p>Indoor:</p> <p>Calcium without D3: each meal</p> <p>Pollen: each meal</p> <p>Multivitamin mix: biweekly</p> <p>Calcium with D3: biweekly</p> <p>Outdoor:</p> <p>Calcium without D3: each meal</p> <p>Pollen: each meal</p> <p>Multivitamin mix: biweekly</p>
	Hydration	<p>Hydration is to be facilitated by combination of night fogging, morning and evening misting and daytime dripping.</p> <p>Urates to be assessed and in case of deviations, hydration methods to be adjusted</p>